

## **Assessment of Nitrogen and Phosphorous Applications for Yield Potential of Guar Strains Under Irrigated Conditions**

*Muhammad Azeem Ur Rahman Khalid Khalid<sup>1</sup>*

*Agricultural Research Station, Bahawalpur, Pakistan*

### **Abstract**

The productive efficiency of two guar varieties BR-2017 and BR-99 was studied with variable nitrogen and phosphorous levels including 0-0, 0-60, 15-60, 30-0, 30-30, 30-60, 30-90 and 45-60 NP kg ha<sup>-1</sup> during two sowing seasons 2015 and 2016 at Agriculture Research Station, Bahawalpur. The results showed that BR-2017 performed better than the BR-99. Variable levels of nitrogen and phosphorous also enhanced the yield of guar. The results indicated that maximum average grain yield of guar 2544 kg ha<sup>-1</sup> was recorded in BR-2017 with NP dosage of 30:60 kg ha<sup>-1</sup> nitrogen and phosphorous that was non-significantly different (2457.5 kg ha<sup>-1</sup>) from NP level of 30:90 kg ha<sup>-1</sup> recorded from same variety. Results indicated that 'BR-2017' was more receptive to fertilizer application than the BR-99.

**Keywords:** *Cluster Bean, Nitrogen, Phosphorous, Varieties, Split Plot Design, ANOVA, LSD.*

---

<sup>1</sup>Agricultural Research Station, Bahawalpur, Plant Breeding and Genetics, Pakistan.

**Correspondence:** aurkhalid75@yahoo.com

## Introduction

Cluster bean scientifically known as [*Cyamopsis tetragonoloba* (L.) Taub.] is self-pollinated crop belonging to the family *Leguminosae*. Guar is a potential summer crop having the ability to fix nitrogen from atmosphere and resist under drought conditions because of it have tap root system that goes down and results in more efficacy of water intake (Farencois et al., 1990). Cluster bean is the main crop in drought prone areas of United States, South Africa, India and Pakistan (Ashraf et al., 2005). It performs well under warm climate and with supplementation of irrigation (Alexander et al., 1988) and fertilization (Omer et al., 1993). Guar has the ability to fix nitrogen in the atmospheric due to nodules (Elsheikh and Ibrahim, 1999; Wetselaar, 1967) and can tolerate saline soils (Farencois et al., 1990; Ashraf et al., 2005).

The major portions of arid areas of the Indo-Pak are occupied by guar due to its legume capability. Guar seed have guar gum (galactomannans, 25-35%), 4% edible oil (Mehta and Ramakrishanan, 1957) and 27 to 37% protein (Whistler and Hymowitz, 1979) that are used for industrial purposes like stabilizer, food industry, cosmetics, paint, textile and detergents (Farencois et al., 1990; Jukanti et al., 2015). Guar gum is also used in several industries like mining, sauces, dairy products, meat blinder etc. (Morris, 2004; Kays et al., 2006) as well as beverages, photography, petroleum and paper (Whistle and Hymowitz 1979; Undersander et al., 1991; Pathak et al., 2010). Guar can be used for agricultural purposes like fodder for animals, vegetable green manuring, (Douglas, 2005). Due to its reduced transpiration and deeper root system it can reduce the shortage of forage in the months of deficit of forage (Paleg and Aspinal, 1981).

The cultivation of legumes is limited to neglected lands without application of any artificial fertilizer. The atmospheric nitrogen is fixed by leguminous crops through symbiotic relationship with bacteria to meet their nitrogen requirements (Herridge et al., 1993). The phosphorus and potassium requirement of leguminous crops is more than cereals because NP is required for BNF bacteria and legume crop itself for its better growth. The available potassium (NFDC, 1994) and phosphorus are deficient in most Pakistani soil.(Fatima et al., 2006) and. The application of phosphorus in soil not only enhance the weight per plant (Patel and Kotecha, 2006; Tariq et al., 2007) but also increases its percentage of protein (Joshi and Mali, 2004). The reduced activity of nitrogen fixing bacteria reduced the nitrogen available in the root zone due the deficiency of phosphorus in leguminous crops (Giller and Cadisch, 1995; Rahman et al., 2008). The weight of individual nodule increased upto four times due to the application of phosphorous and potassium. (Lynda et al., 1984).

The inoculation of seed with Rhizobium and NP application significantly affected the grain yield of guar except germination, primary branches, No. of Pods per plant and harvest index (Khan et al., 1992) The application of nitrogen @ 100 kg ha<sup>-1</sup> significantly increase the seed yield and dry weight per plant and inoculation of seed did not affect the seed yield (McKenzie et al., 1992). The application of nitrogen @ 25 kg ha<sup>-1</sup> enhanced the grain yield of chickpea (1.3 ton ha<sup>-1</sup>) (Kasole et al., 1995). The application of

nitrogen @ 25 kg ha<sup>-1</sup> enhanced the protein contents and seed yield however seed yield and protein contents was stable up to 50 kg N per ha (Kurahde *et al.*, 1994). Therefore, nitrogen and phosphorus are essential for better growth of crop and nutritional status in dry matter. The present study was conducted to assess the growth and yield of guar cultivars BR-99 and BR-2017 to variable NP doses under irrigated conditions of Bahawalpur.

### **Material and Methods**

In this study, two varieties of Guar viz; BR-99 (V1) and BR-2017 (V2) were grown with eight different levels of NP fertilizer at the Agricultural Research Station, Bahawalpur during 2015 and 2016. The fertilizer levels T1 (0-0), T2 (0-60), T3 (15-60), T4 (30-0), T5 (30-30), T6 (30-60), T7 (30-90) and T8 (45-60) kg NP ha<sup>-1</sup> were applied at the time of land preparation before sowing. The trial was laid out in a split plot design with three replications. Varieties were kept in main plot and fertilizer levels in sub-plots. A plot size of 1.8 m x 6 m keeping six (4) rows 45 cm apart was maintained. The whole quantity of nitrogen and phosphorus was applied in the form of urea and single super phosphate, respectively. All other cultural practices were kept uniform in all the plots. At physiological maturity of the crop, yield data were recorded. The samples harvested for grain yield were sun dried and threshed. The data were subjected to ANOVA and the treatments means were compared by LSD at 5% level of probability (Steel & Terrie, 1984).

### **Results and Discussions**

The highest average seed yield of guar 2544 kg ha<sup>-1</sup> was obtained by BR-2017 with the application of 30-60 kg ha<sup>-1</sup> NP that was statistically at par with 30-90 kg ha<sup>-1</sup> NP fertilizer (2457.5 kg ha<sup>-1</sup>). The results indicated that 'BR-2017' variety performed well to the application of NP fertilizer as compared to BR-99. The minimum average seed yield of guar (842 kg ha<sup>-1</sup>) was recorded in BR-99 variety in control where no fertilizer was applied (Table 1).

Treatments, varieties and treatments × varieties interactions were highly-significant and having a difference between all varieties which showed different yield potential at different level of fertilizers. There were no significant differences between blocks (Table 2, 5).

Fertilizer level T6 indicated maximum yield potential. The data showed that av. grain yield of guar at variable NP levels was between 968.66 kg ha<sup>-1</sup> to 2144.5 kg ha<sup>-1</sup>. The average grain yield was non-significant with the appliance of 30-90 and 45-60 kg NP ha<sup>-1</sup>. The optimal level for maximum production of grain yield of guar is 30-60 kg NP ha<sup>-1</sup> because the inadequate application of fertilizer results vegetative growth resulting in loss of grain yield (Table 3, 6).

BR-2017 showed maximum yield potential than BR-99. Average grain yield data indicated that both the guar varieties responded well to variable levels of NP and varieties differ significantly among treatment

means. The results on average grain yield of guar varieties indicated that both the varieties differ significantly to various levels of NP. The yield parameters like no of pod plant<sup>-1</sup> and no of pods cluster<sup>-1</sup> of BR-2017 and BR-99 also supported the increase in yield of guar. (Table 4, 7).

The grain yield of both cultivars BR-2017 and BR-99 was significantly increases with variable levels of nitrogen and phosphorous. The above discussed consequences are comparable with Yaqoob et al. (2005) and Bashir et al. (2006) i.e. chickpea and lentil varieties differ among yield that are developed through selection. Thus, the 'BR-99' is recommended for cultivation for the purpose of increase in guar yield compared to other varieties of Guar.

**Table 1.** Average yield data of guar varieties kg ha<sup>-1</sup> at different levels of NP 2015 and 2016.

Year		2015		2016	
Variety		BR-99	BR-2017	BR-99	BR-2017
<b>T1</b>	0-0	859.5	1115.5	842.0	1089.0
<b>T2</b>	0-60	1167.5	1181.0	1512.0	1644.0
<b>T3</b>	15-60	1221.0	1507.5	1318.0	2182.5
<b>T4</b>	30-0	1061.5	1366.5	982.0	1370.5
<b>T5</b>	30-30	1484.0	1970.0	1646.0	2331.5
<b>T6</b>	30-60	1694.0	2366.0	1797.0	2544.0
<b>T7</b>	30-90	1690.5	2212.0	1689.0	2457.5
<b>T8</b>	45-60	1482.5	2095.0	1573.0	2058.5

**Table 2.** ANOVA Of yield data of two guar varieties at various NP levels during 2015.

ANOVA	D.F	SS	MS	F cal	F tab
Block	2	78327.1	39163.5	1.8	0.3 NS
Varieties	1	2252466.7	2252466.7	106.5	0.009 **
Error 1	2	42274.5	21137.2		
Treatment	7	7749240.2	1107034.3	36.4	0.0 ***
V × T	7	326824.9	46689.2	7.5	0.2 **
Error 2	28	849728.3	30347.4		
Total	47	11298861.9			

**Table 3.** Yield ranking of guar by the effect of N and P fertilizers 2015.

Rank Mean	Level NP	Mean(kg ha <sup>-1</sup> )	Ranking
T6	30-60	2045.6	A
T7	30-90	1957.1	AB
T8	45-60	1848.3	AB
T5	30-30	1692.3	B
T3	15-60	1355.6	C
T4	30-0	1204.6	CD
T2	0-60	1013.1	DE
T1	0-0	968.6	E

**Table 4.** Grain yield of guar varieties as influenced by fertilizer application 2015

Variety		Mean(kg ha <sup>-1</sup> )	Ranking
BR-2017	V2	1727.33	A
BR-99	V1	1294.083	B

**Table 5.** ANOVA of yield data of two guar varieties at various NP levels during 2016

ANOVA	D.F	SS	MS	F cal	F tab
Block	2	195102.1	97551.062	0.9	.50NS
Variety	1	3441123	3441123	34.2	.02 *
Errore1	2	200765.3	100382.6		
Treatment	7	7236619.2	1033802.8	66.7	.0 ***
V × T	7	1081801.6	154543.1	9.9	.0 ***
Errore2	28	433429.8	15479.6		
Total	47	12588841.2			

**Table 6.** Yield ranking of guar by the effect of N and P fertilizers 2016.

Rank Mean	Level NP	Mean (kg ha <sup>-1</sup> )	Ranking
T6	30-60	2144.5	A
T7	30-90	2122.3	B
T5	30-30	1912.6	B
T8	45-60	1900.8	B
T3	15-60	1734.6	C
T2	0-60	1499.8	D
T4	30-0	1199.1	E
T1	0-0	1027	F

**Table 7.** Grain yield of guar varieties as influenced by fertilizer application 2016.

Variety		Mean (kg ha <sup>-1</sup> )	Ranking
BR-2017	V2	1960.3	A
BR-99	V1	1424.8	B

### Conclusions

Both the varieties differed significantly for yield potential by utilization of different levels of fertilizer. An improved variety “BR-2017” responded more efficiently to fertilizer than BR-99 variety. Thus, it can be concluded that the farmers can adopt newly approved variety of guar with appropriate doses of fertilizer (30-60 kg NP ha<sup>-1</sup>) application to increase their yield.

### References

- Ashraf, M. Y., K. Akhtar, G. Sarwar and M. Ashraf. (2005). Role of rooting system in salt tolerant potential of different guar accessions. *Agron. Sustain. Dev.*, 25: 243-249
- Alexandar, W. L., D. A. Buck and R. A. Backhaus. (1988). Irrigation water management of guar seed production. *Agron. J.*, 80: 447- 453.
- Ali, L., A.R. Chaudry and A.H. Shah. (1984). Correlation between yield and yield component in wheat. *J. Agric. Res.*, 22: 279–83.
- Bashir, A., M. Yaqoob and M. Rahim. (2006). Karak-2, a new high yielding variety of chickpea for rain fed ecologies of NWFP. *Indus J. Pl. Sci.*, 5: 639–41.
- Douglas, C. A. (2005). Evaluation of guar cultivars in central and southern queensland. *A report for the rural industries research and development corporation*. Pp:3.
- Elsheikh, E. A. E. and K. A. Ibrahim. (1999). The effect of Bradirhizobium inoculants on yield and seed quality of guar (*Cyamopsis tetragonoloba* L.). *Food Chemistry*, 65: 183-187.
- Farencois, L. E., T. J. Donovan and E. V. Maas. (1990). Salinity effects on emergence, vegetative growth and seed yield of guar. *Agron. J.*, 82: 587-592.
- Fatima, Z., M. Zia and M. F. Chaudhary. (2006). Effect of rhizobium strains and phosphorus on growth of soybean (*Glycine max*) and survival of *rhizobium* and P solubilizing bacteria. *Pak. J. Bot.* 38: 459- 464.
- Joshi, N. L. and P. C. Mali. (2004). Effect of application of phosphorus and time of harvest on seed, crude protein and gum yield of cluster bean. *J. Arid Legumes*. 1:27-3.

- Jukanti, A., R. Bhatt, R. Sharma and R. Kalia. (2015). Morphological, agronomic, and yield characterization of cluster bean (*Cyamopsis tetragonoloba* L.) germplasm accessions. *J. Crop Sci. Biotechnol.*, 18: 83-88.
- Giller, K. E. and G. Cadisch. (1995). Future benefits from biological nitrogen fixation: An ecological approach to agriculture. *Plant Soil*. 174: 255- 277.
- Herridge, D. F., O. P. Rupela, R. Sarraj and D. P. Beck. (1993). Screening techniques and improved biological nitrogen fixation in cool season legumes. *Euphytica*. 1: 1-14.
- Kasole. K.E., S.D. Kalke, S.M. Kareppa and K.K. Khade. (1995). Response of chickpea (*Cicer arietium* L.) to different fertilizer levels. plant population and weed management on cultivator's field in north eastern parts of Kolhapur. Maharashtra. *Indian J. Agron.* 40: 217-219.
- Kays, S. E., J. B. Morris and Y. Kim. (2006). Total and soluble dietary fiber variation in [*Cyamopsis tetragonoloba* (L.) Taub.] (Guar) genotypes. *J. Food Quality*, 29:383–391
- Khan, H. R., A. M. Haqqani, M. A. Khan and B. A. Malik. (1992). Biological and chemical fertilizer studies in chickpea grown under arid conditions of Thai. *Sarhad J. Agri.* 8: 321-327.
- Kurhade. N. G., L. A. Deshpande and K. T. Nagre. (1994). Effect of nitrogen and phosphorus on the yield and quality of gram cultivars. *J. Maharashtra Agri. Univ.* 19: 277-278.
- Lynda, J. Q., E. A. H. Jra and G. V. O. Jra. (1984). Nodulation and nitrogen fixation by arrow leaf clover: Effects of phosphorus and potassium. *Soil Biol. Bioch.* 16: 589-594.
- Mehta, D. R. and C. V. Ramakrishanan. (1957). Studies of guar seed (*Cyamopsis psoralioides*). *Journal of American Oil Chemistry Society*, 34: 459-461.
- McKenzie, B. A., M. Andrews. A. Z. Ayalsc« and J. R. Stokes. (1992). Leaf growth and canopy development in chickpea. *Proc. Annual Conference. Agronomy Society or New Zealand.* 22: 121-125.
- Morris, J. B. (2004). Legumes: nutraceutical and pharmaceutical uses. In: Goodman RM (ed) *Encyclopedia of plant and crop science*. Marcel Dekker, New York, pp 651–655.
- NFDC. (1994). Sulphur status and crop response in Pakistan soils. *NDFC publication No.2/94*, NDFC, Islamabad.
- Rahman, M. M., M. M. H. Bhuiyan, G. N. C. Sutradhar, M. M. Rahman and A. K. Paul. (2008). Effect of phosphorus, molybdenum and rhizobium inoculation on yield and yield attributes of mungbean. *Int. J. Sustain. Crop Prod.* 3:26-33.

- Shah, S. H., D. F. Khan and M. S. Madaui. (1994). Effect of different rhizobial strains on the performance of two chickpea cultivars under field conditions. *Sarhad J. Agric.* 10: 103-107.
- Steel, R. G. D. and I. H. Torrie. (1984). *Principles and Procedures of Statistics*. 2nd ed. McGraw Hill Book Co Inc. Singapore: 172-177.
- Tariq, S., S. Ali and S. S. Ijaz. (2007). Improving nitrogen fixation capacity and yield on mung bean and mash bean by phosphorous management in pothowar. *Sarhad J. Agric.* 23:1027-1032.
- Omer, E. A., A. Fattah, M. Razin and S. S. Ahmed. (1993). Effects of cutting, phosphorus and potassium on guar (*Cyamopsis tetragonoloba*) in newly claimed soil in Egyptian. *Plant Food for Human Nutrition*, 47: 277-284.
- Paleg, L. G. and D. Aspinal. (1981). *The physiology and biochemistry of drought resistance in plants*. Academic press, Sydney, Australia. pp. 251-252
- Patel, P. C. and A. V. Kotecha. (2006). Effect of phosphorus and potassium on growth characters, forage yield, nutrient uptake and quality of Lucerne (*Medicago sativa* L.). *Ind. J. Agron.* 51: 242-244.
- Pathak, R., S. K. Singh, M. Singh and A. Henry. (2010). Molecular assessment of genetic diversity in cluster bean (*Cyamopsis tetragonoloba*) genotypes. *J. Genetics.*, 89: 243-246.
- Undersander, D. J., D. H. Putnam, A. R. Kaminski, K. A. Kelling, J. D. Doll, E. S. Oplinger and J. L. Gunsolus. (1991). Guar. In: *Alternative Field Crops Manual*. Uni. of Wisconsin-Extension. <http://www.hort.purdue.edu/newcrop/afcm/guar.html>.
- Wetselaar, R. (1967). Estimate of nitrogen fixation by four legumes in a dry monsoonal area north-western Australia. *Australian Journal of Experimental Agriculture and Animal Husbandry*, 7: 518-522.
- Whistler, R. I. and T. Hymowitz. (1979). *Guar: agronomy, production, industrial use, and nutrition*. Purdue University Press, West Lafayette, Indiana, USA.
- Yaqoob, M., M. Mansoor, Najibullah and Nasirudin, (2005). A high yielding lentil cultivar for rainfed ecologies of NWFP. *Indus J. Pl. Sci.*, 4: 314-8